arXiv:math/0411106v1 [math.MG] 4 Nov 2004

Hypercubes circumscribed in hyperspheres: a constant growth ratio for volumes at large dimensions

N. Shabot-Marcos[a] y A. Sandoval-Villalbazo[a]
[a] Departamento de Física y Matemáticas, Universidad Iberoamericana
Lomas de Santa Fe 01210 México D.F., México
E-Mail: alfredo.sandoval@uia.mx
E-Mail: nathanshabot@alumno.uia.mx

February 1, 2008

Abstract

This note shows that an interesting property arises when considering the relation between the hypersphere volumes at dimensions $n+1$ and n, if the hyperspheres circumscribe unitary hypercubes in $n+1$ and n dimensions, respectively . In the limit of large n, the ratio $\frac{V_{n+1}}{V_n}$ is constant and equal to the irrational number $\sqrt{\frac{\pi e}{2}}$. Some consequences of this result are explored.

It is well known that the volume of a hypersphere embedded in an euclidean n-dimensional space decreases to zero for large n [1] [2]. The value of n in which the hypersphere volume starts to decrease depends on the ratio R. On the other hand, the corresponding volume of a unitary hypercube remains constant in an euclidean n-dimensional space. We now consider the problem of the circumscription of hypercubes in hyperspheres in euclidean n-dimensional spaces.

The volume of a hypersphere in n dimensions, for a given ratio R, is given by:

$$V_n = \frac{2\pi R^n}{n} \prod_{k=1}^{n-2} \frac{\sqrt{\pi}\ \Gamma(\frac{k+1}{2})}{\Gamma(1+\frac{k}{2})},\ n \geq 3. \tag{1}$$

If the hypersphere circumscribes an unitary hypercube in the same n dimensional space, then, the hypersphere ratio is given by:

$$R = \frac{\sqrt{n}}{2} \tag{2}$$

So, the volumes V_{n+1} and V_n are related by the equation:

$$g_n = \frac{V_{n+1}}{V_n} = \frac{n^{\frac{n}{2}-1}}{2}(n-1)^{\frac{3-n}{2}} \frac{\sqrt{\pi}\ \Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \tag{3}$$



Figure 1: The expression $\frac{V_{n+1}}{V_n}$ for the case of hyperspheres of dimensions $n+1$ and n circumscribing unit hypercubes of the respective dimensions.

Clearly, for large n:

$$\lim_{n\to\infty} g_n = \sqrt{\frac{\pi e}{2}} \simeq 2.0663656 \tag{4}$$

Fig. 1 shows, for continuous n, the corresponding plot of Eq.(3)

To the authors' knowledge, this is the first time that the irrational number $\sqrt{\frac{\pi e}{2}}$ appears in a computation related to n-dimensional geometry. The result shows that the circumscription requirement defines o family of hyperspheres that possess constant volume growth as the number of dimensions of its corresponding space tends to infinity. Further properties of circumscription problems will be studied in future work.

References

[1] Sommerville, D. M. Y. An Introduction to the Geometry of n Dimensions. New York: Dover, p. 136, 1958.

[2] Eric W. Weisstein. "Hypersphere." From MathWorld–A Wolfram Web Resource. http://mathworld.wolfram.com/Hypersphere.html, and references therein.